

A RAND NOTE

How Coronary Angiography Is Used: Clinical Determinants of Appropriateness

**Mark R. Chassin, Jacqueline Kosecoff,
David H. Solomon, Robert H. Brook**

November 1987

RAND

The research described in this report was sponsored by the Commonwealth Fund, the John A. Hartford Foundation, the U.S. Health Care Financing Administration, U.S. Department of Health and Human Services, the Pew Memorial Trust, and the Robert Wood Johnson Foundation.

This Note contains an offprint of RAND research originally published in a journal or book. The text is reproduced here, with permission of the original publisher.

The RAND Publication Series: The Report is the principal publication documenting and transmitting RAND's major research findings and final research results. The RAND Note reports other outputs of sponsored research for general distribution. Publications of The RAND Corporation do not necessarily reflect the opinions or policies of the sponsors of RAND research.

A RAND NOTE

N-2749-CWF/HF/HCFA/PMT/RWJ

**How Coronary Angiography Is Used:
Clinical Determinants of Appropriateness**

**Mark R. Chassin, Jacqueline Kosecoff,
David H. Solomon, Robert H. Brook**

November 1987

**Prepared for
The Commonwealth Fund
The John A. Hartford Foundation
The U.S. Health Care Financing Administration,
U.S. Department of Health and Human Services
The Pew Memorial Trust
The Robert Wood Johnson Foundation**

RAND

How Coronary Angiography Is Used

Clinical Determinants of Appropriateness

Mark R. Chassin, MD, MPP, MPH; Jacqueline Kosecoff, PhD;
David H. Solomon, MD; Robert H. Brook, MD, ScD

Using ratings of appropriateness derived from an expert physician panel, we measured how appropriately physicians in 1981 performed coronary angiography in a randomly selected, community-based sample of cases in the Medicare population. We studied large geographic areas (three sites) in three states, representing regions of high and low use. The high-use site had fewer procedures classified as appropriate (72%) than either low-use site (77% and 81%, respectively). Over all sites, 17% of procedures were classified as inappropriate. Patients in the high-use site were older, had less severe angina, and were less intensively medically treated than patients in either of the low-use sites. Patients without angina who had not undergone exercise testing constituted the most common subgroup of inappropriate cases. Although overall differences in appropriateness were not large, practice differences do exist. This analysis of practice differences among study sites provides the clinical basis for understanding the small, but significant, differences in the appropriateness of use of coronary angiography. The finding of 17% inappropriate use may be cause for concern.

(*JAMA* 1987;258:2543-2547)

CORONARY angiography is an invaluable diagnostic tool; no other available technique can delineate the anatomy of the coronary circulation as precisely. Yet it is also an expensive procedure, with rates of major complications that are low but clinically important.^{1,6}

Substantial differences exist from one geographic area to another in the population-based rates at which this

procedure is performed.⁷ Questions have been raised concerning the extent to which it is used inappropriately.⁸ While many studies have contributed to our knowledge about the circumstances under which coronary angiography is

See also pp 2533, 2538, and 2568.

efficacious, we know little about how the procedure is actually used in community practice.

We measured the appropriateness with which coronary angiography was performed in the elderly Medicare population of three large geographic areas in 1981. In the first article in this series, we have reported the relationship between appropriateness and rates of use in these three areas.⁹ We report herein the clinical characteristics of the patients who underwent coronary an-

giography, the importance of these clinical factors in determining appropriateness, and how they differed among the three geographic areas.

Methods

We used Medicare physician claims data to measure population-based rates of use in 1981 of coronary angiography and other health care services in 13 large geographic areas across the United States. The methods and results of this analysis have been reported previously.⁷ The mean rate of coronary angiography among these 13 sites in 1981 was 33 per 10 000 (adjusted for age and sex differences). The rates we observed ranged from a low of 22 to a high of 51 per 10 000; the differences among the 13 sites were statistically significant ($P < .001$, χ^2).

We selected three sites with rates at both extremes of the distribution in which to carry out the study of appropriateness. Site 1 had a high-use rate, and is a mixed urban and rural area. We selected two low-use sites for study because these two sites have contrasting demographic characteristics; site 3 is relatively rural, while site 2 is more urban.

We have reported the methods by which we reviewed the medical literature, developed a catalog of indications for coronary angiography, and convened a panel of expert physicians to rate each indication on a nine-point scale of appropriateness.^{6,9-11} The panel was composed of three cardiologists, two internists, one family physician, two cardiac surgeons, and one radiologist. They rated 300 separate indications for

From the Health Program of The Rand Corp (Drs Chassin and Brook), and Fink and Kosecoff Inc (Dr Kosecoff), Santa Monica, Calif; and the Departments of Medicine (Drs Kosecoff, Solomon, and Brook), and Public Health (Drs Kosecoff and Brook), University of California at Los Angeles.

The opinions, conclusions, and proposals in the text are those of the authors alone and do not necessarily represent the views of any of these organizations, The Rand Corp, or the University of California.

Reprint requests to The Rand Corp, 1700 Main St, PO Box 2138, Santa Monica, CA 90406-2138 (Dr Chassin).

coronary angiography.⁶ (Rand reports are available at nominal cost on request from The Rand Corporation, Santa Monica, Calif.) The definition of appropriateness the panel used, the method by which it was conducted, and the way indications were classified as appropriate, equivocal, and inappropriate are described in the first article in this series.⁹ In general, we used median ratings to classify indications as appropriate (ratings of 7 to 9), equivocal (ratings of 4 to 6), or inappropriate (ratings of 1 to 3). Where our panelists disagreed on appropriateness ratings, however, we classified such indications as equivocal, regardless of median rating.

Using the claims data from which we calculated overall rates of use, we selected a random sample of cases from each of the three sites. We have described in the preceding article in this series how we obtained the participation of the community-based sample of physicians who performed these coronary angiograms.¹² The overall physician participation rate for coronary angiography was 91%, comprising some 219 physicians. There were no significant differences among sites in the rates of physician participation (89%, site 1; 96%, site 2; and 93%, site 3). Participating physicians had performed 95% of sampled procedures (N = 1810).

We sought the medical records of all 1810 cases represented by participating physicians. We developed a detailed abstraction form to collect the data necessary to assign indications to each case. For example, we collected data on the nature of any chest pain (its quality, location, severity, provoking factors, and stability); results of exercise tests (treadmill, thallium, or gated-blood pool scan); medication schedules; occurrence, dates, sites, and complications of prior myocardial infarctions; and occurrence of previous coronary artery bypass graft surgery and which vessels were bypassed. We obtained photocopies of the results of all exercise tests and of the coronary angiography reports of all studied procedures. The abstraction form and guidelines for its use have been published elsewhere.⁶

Our panel used the Canadian Cardiovascular Society classification to grade the severity of angina. They defined maximal medical therapy as the use of any long-acting nitrate and any β -blocker (or a documented history of intolerance to either). We defined angina as present if any three of the following four conditions were met: chest pain present in typical location (eg, substernal, left side of the chest), pain of typical quality (eg, pressing or tight), pain typically produced by exer-

tion, and pain typically relieved by sublingual nitroglycerin therapy. A positive exercise treadmill test result was defined as 1 mm or more of horizontal or down-sloping ST-segment depression, or the occurrence of typical angina at any time during the test after the first three minutes. A very positive exercise treadmill test result was defined as a positive test result occurring during the first three minutes of exercise, or a drop in systolic blood pressure of 20 mm Hg or more occurring during the first three minutes of exercise, or more than 2.5 mm of horizontal or down-sloping ST-segment depression at any time.

Data collection took place in hospitals, physicians' offices, and clinics as necessary to obtain complete clinical data on each case. Data collection methods are described in the preceding article in this series.¹² Of the 1810 procedures of participating physicians sampled from the claims data, we determined that 75 (4.1%) represented errors in the claims data: the procedure performed was not a coronary angiography (1.9%), the procedure was not performed in 1981 (0.4%), the procedure was not performed within the geographic area (1.8%), or the physician had no record at all of having seen the patient (1.7%).

We were able to obtain complete information in 1677 cases, or 97% of all cases eligible for inclusion in the study after excluding the claims errors. Data were incomplete in 58 cases (3%) and were excluded either because medical records known to exist could not be found (56 cases) or because available medical records from all sources could not provide data complete enough for us to determine the indication for the procedure (two cases).

Based on data elements recorded in the abstraction forms, one or more indications was assigned to each case. For cases with more than one indication, the one with the highest appropriateness rating was designated the primary indication and was used throughout the subsequent analysis. In the preceding article in this series, we reported studies of the validity of the complete process of medical records abstraction and indications assignment.¹²

We sampled roughly equal numbers of cases in each site to maximize our ability to detect differences among sites. With 500 cases in each site, we could expect to detect a true difference of eight percentage points in appropriate or inappropriate use in at least 80% of repeated trials. We tested for differences in the distributions among sites of patient characteristics and indications using χ^2 tests.¹³

RESULTS

Demographic and clinical characteristics of the patient populations in each site are listed in Table 1. The sites differ on several of these factors. Patients in the high-use site were significantly older than in either of the low-use sites; 21% of patients in site 1 were older than 74 years, compared with 15% in site 2, and 9% in site 3. Patients in the high-use site were also more likely to have had a previous coronary angiogram or previous coronary artery bypass surgery (CABS) than in either of the low-use sites.

Significant differences among sites were observed in the proportion of patients with previous myocardial infarction and those with congestive heart failure. These differences were not, however, consistently related to the rate of use of coronary angiography. No significant differences among sites were observed in race or sex or the proportion of patients with angina.

We divided the 300 indications for coronary angiography into 12 major clinical groups according to the patient's major presenting complaint. The distribution of cases whose primary indications fell in each major clinical group in each site is shown in Table 2. In general, patients received coronary angiography for the same reasons in all sites. In site 1, chronic stable angina, unstable angina, and valvular disease accounted for 64% of patients; in site 2, for 66%; and in site 3, for 65%.

Only minor differences among sites are apparent in the distributions of patients by major clinical group. Somewhat fewer patients in the high-use site received their coronary angiograms during or after hospital admission for unstable angina than in the low-use sites: 22% in site 1 compared with 29% in site 2, and 28% in site 3. As suggested by data in Table 1, more patients in site 1 had their coronary angiograms following CABS than in either of the low-use sites. Other differences were minor and inconsistently related to overall rates of use.

Table 3 shows how the cases in each site were distributed by groupings of indications for each appropriateness category and gives more detail about severity of symptoms, results of noninvasive tests, and level of medical therapy. The high-use site (site 1) demonstrated significantly fewer appropriate cases than either of the low-use sites. The high-use site showed more equivocal cases than either of the low-use sites, but there were no significant differences in the distribution of inappropriate cases by site.

Table 1.—Demographic and Clinical Characteristics of Patients Undergoing Coronary Angiography in Three Geographic Areas in 1981

Characteristic*	Site No.		
	1 (n=628)	2 (n=514)	3 (n=535)
Coronary angiography rate/10000 elderly†	50	23	22
Age, y, %‡			
64-69	47	55	63
70-74	31	31	27
75-79	15	13	8
≥80	6	2	1
Sex, %§			
M	60	58	58
Nonwhite, %§	6	9	7
Congestive heart failure, %	21	27	20
Angina, %§	37	36	37
Previous MI, %¶			
0	55	47	63
1	35	37	28
>1	10	16	9
Previous CABS, %‡	9	4	3
Previous coronary angiography, %‡	15	9	8

*MI indicates myocardial infarction; and CABS, coronary artery bypass surgery.
 †The rates for persons aged 65 years or older in 1981 are adjusted for differences in age and sex among sites.
 ‡ $P < .01$ for differences in age distribution among sites.
 § $P > .05$ for differences among sites in sex and race distributions and in percent of patients with angina.
 || $P < .05$ for differences among sites in percent of patients with congestive heart failure.
 ¶ $P < .01$ for differences among sites in percent of patients with none, one, or more than one previous MI.

To understand better the differences among sites in distribution of cases by appropriateness category, we examined the severity of patients' angina and degree of medical therapy in each site. Data in Table 4 show that for patients with clinical manifestations of coronary artery disease, those in site 1 had significantly less severe angina than patients in either low-use site. Patients in site 1 were also less frequently treated with long-acting nitrates and β -blockers, a regimen our panel defined as maximal medical therapy for 1981. In site 1, of the patients with angina, 52% received maximal therapy compared with 70% in site 2, and 63% in site 3 ($P < .01$, χ^2).

We looked further at intensity of medical management by measuring the daily dose of propranolol hydrochloride received by each patient with angina receiving this drug. In our sample of 1981 patients, this drug was by far the most frequently used β -blocker. Among all sites, 81% of patients receiving β -blockers were taking propranolol (81%, site 1; 83%, site 2; and 78%, site 3; $P > .05$, χ^2). Data in Table 4 show the distribution of patients according to their daily dose of propranolol in each site. More patients with angina receiving propranolol received low daily doses in site 1 than in either low-use site.

Finally, data in Table 4 also show the results of angiography by site for those patients with angina (stable or unsta-

Table 2.—Distribution of Patients Who Underwent Coronary Angiography by Major Clinical Group and Site

Major Clinical Group, %*	Site No.†		
	1 (n=628)	2 (n=514)	3 (n=535)
Chronic stable angina	28	25	28
Unstable angina‡	17	22	20
Valve disease	19	19	17
Following MI§	10	12	9
Nonspecific chest pain	8	4	11
Following unstable angina	5	7	8
Following CABS	5	2	1
Asymptomatic	3	1	2
During an acute MI¶	2	2	1
Congestive heart failure	2	3	2
Following survival from sudden cardiac arrest	<1	<1	<1
Other	1	1	2

*MI indicates myocardial infarction; and CABS, coronary artery bypass surgery.
 †Column percentages may not add up to 100% due to rounding.
 ‡Coronary angiography was performed during the hospital admission occasioned by unstable angina.
 §This is defined as within six months of an MI (but more than six days after its onset).
 ||This is defined as within three months of hospital discharge following an episode of unstable angina.
 ¶This is defined as within six days of the onset of MI.

ble) or other clinical manifestations of coronary artery disease (myocardial infarction or previous CABS). There were no significant differences among sites in the severity of coronary artery disease discovered at the time of angiography. The results were the same whether we used a criterion of 50% reduction in luminal diameter or the 70% criterion used in the data presented in Table 4.

COMMENT

We studied how appropriately physicians used coronary angiography in 1981 in the elderly Medicare population. We examined cases from three large geographic areas, one with a high rate of use and two with low rates of use for this procedure. The fact that patients in the high-use site were more likely to have had a previous coronary angiogram or CABS suggests that these patterns of high- and low-use were present for some time before 1981.

We found that physicians used coronary angiography differently in the high-use site compared with their counterparts in the two low-use sites. Patients in site 1 were older, had less severe angina, and were less often treated with maximal medical therapy than patients in the low-use sites. These findings help explain our finding of fewer appropriate and more equivocal procedures in site 1 than sites 2 or 3. Over all sites, about three fourths of the cases were classified appropriate.

There were no significant differences among sites in the frequency of cases classified as inappropriate. There were, however, a substantial proportion of inappropriate procedures in all three sites, averaging 17% of all cases. Patients who had no angina and no func-

tional evaluation by means of exercise testing constituted the most frequent group, accounting for about half of all inappropriate cases over all three sites. Among these patients, about two thirds had previously sustained a myocardial infarction, undergone CABS, had stable angina, or been hospitalized for unstable angina. The remaining one third were entirely asymptomatic or had nonspecific chest pain.

The results of coronary angiography in our patients showed no differences in severity of coronary artery disease among sites. We expected to find somewhat less severe disease in site 1, because patients there had less severe symptoms and less intensive medical treatment than patients in the other sites. The most likely explanation for this finding is that patients in site 1 were significantly older than their counterparts in the other sites. Probably the effect of increased age cancelled out the effects of decreased levels of symptoms and treatment.

The principal factor that might limit the generalizability of our results is that we studied the use of coronary angiography in 1981. Our expert panel rated the appropriateness of performing this procedure based on information known to have been available and known standards of practice in 1981. What has changed since 1981? The most important changes have been the addition of new data on the efficacy of CABS and the emergence of several new treatments: calcium-channel blocking agents for the medical treatment of angina, intracoronary thrombolytic treatment of acute myocardial infarction, and transluminal angioplasty as an alternative to CABS.

Table 3.—Indications for Coronary Angiography by Site for Each Appropriateness Category

Indications, %*	Site No.		
	1 (n = 628)	2 (n = 514)	3 (n = 535)
Appropriate			
During evaluation of valve disease	19	19	17
Unstable angina			
Pain controlled	12	13	12
Pain persists	4	8	7
Stable angina			
Class III or IV, maximal therapy	8	12	12
Class III or IV, less than maximal therapy, positive or very positive treadmill test result	3	1	3
Class I or II, maximal therapy, positive or very positive treadmill test result	3	2	2
Class I or II, less than maximal therapy, positive or very positive treadmill test result	4	2	5
Stable angina following MI, unstable angina, or CABS			
Class III or IV	7	14	10
Class I or II	3	2	2
Nonspecific chest pain with positive or very positive treadmill test result	4	1	3
Following MI, unstable angina, or CABS, no angina, with positive or very positive treadmill test result	2	1	4
Other appropriate	3	2	4
Total % appropriate†	72	77	81
Equivocal			
Stable angina			
Class III or IV; less than maximal therapy; no or negative treadmill test result	4	3	2
Class I or II, maximal therapy; no or negative treadmill test result	1	2	1
No angina (asymptomatic, post MI, or previous stable angina), positive treadmill test result; negative or absent exercise thallium scan test result	4	1	1
Other equivocal	2	1	1
Total % equivocal‡	10	7	4
Inappropriate			
No angina, no exercise tests performed, post MI, post CABS, following unstable angina, or previous stable angina	6	5	1
Asymptomatic or nonspecific chest pain	3	3	5
No angina, negative treadmill test result, post MI, post CABS, following unstable angina, or previous stable angina	2	<1	1
Nonspecific chest pain	2	<1	3
Stable angina, class I or II, less than maximal therapy, absent or negative treadmill test result	2	2	1
Congestive heart failure	2	3	2
During acute, uncomplicated MI§	1	1	<1
Other inappropriate	1	1	1
Total % inappropriate	18	17	15

*MI indicates myocardial infarction; and CABS, coronary artery bypass surgery.

† $P < .01$ for differences in frequency of appropriate indications among all sites and for site 1 vs sites 2 and 3 combined.

‡ $P < .01$ for differences in frequency of equivocal indications among all sites and for site 1 vs sites 2 and 3 combined. Column percentages may not add up to total percentages due to rounding.

§These cases represent the use of coronary angiography solely for diagnostic purposes and do not include instances where intracoronary streptokinase was administered; seven such cases were excluded from the analysis.

|| $P > .05$ for differences in frequency of inappropriate indications among all sites and for site 1 vs sites 2 and 3 combined. Column percentages may not add up to total percentages due to rounding.

In 1981, the available data suggested that many, if not all, patients with angina who had three-vessel disease and reasonable ventricular function benefited from CABS, with improved survival.^{6,14-16} Recent studies have docu-

mented that in patients with mild angina this benefit is probably limited to those patients with three-vessel disease who have modest reductions in ejection fraction.¹⁷⁻¹⁹ Medical treatment of angina has improved significantly since

1981, with the addition of calcium-channel blocking agents and a wider spectrum of β -blockers. The increased efficacy of medical treatment,²⁰⁻²² and our improved understanding of the role of CABS might lead experts today to view the use of coronary angiography in patients with mild angina or who have received less than maximal medical therapy as less appropriate than our panel did for 1981.

The use of intracoronary thrombolytic therapy today is limited to a few institutions and not often used in the elderly because of the prevalence of complicating comorbid conditions. Furthermore, it may be supplanted by the intravenous use of tissue plasminogen activator.²³ Because research has yet to document the efficacy of transluminal angioplasty in enhancing survival, its principal appropriate use today is to treat anginal pain. As just discussed, the increased spectrum of medical treatment for angina may have reduced the appropriateness of proceeding to invasive treatment before medical therapy has been exhausted.

In summary, it seems unlikely that any of the major uses of coronary angiography we classified as inappropriate in 1981 would today be considered appropriate. Possibly some of the uses we described as appropriate for 1981—particularly for patients with mild angina and those with less than maximal medical therapy—might be considered as less than appropriate today.

We believe our method of rating the appropriateness of indications for coronary angiography has validity; it separates clinically important differences in appropriateness among different indications for the procedure. While we have not demonstrated predictive validity (eg, that patients who received coronary angiography for indications rated appropriate by our panel have better health outcomes than if they had not received it), we believe the ratings demonstrate face validity and content validity.

Examining the data in Table 3, we believe most physicians will agree that the appropriate indications in the first section of the table represent circumstances under which coronary angiography is clearly indicated, many of which have been well documented in the clinical literature. The appropriateness of coronary angiography for the equivocal indications is far less clear. For the inappropriate indications, we believe few physicians would advocate the routine use of coronary angiography. There is much less, if any, solid research data documenting the efficacy of coronary angiography in these circumstances.

Table 4.—Angina Class, Propranolol Hydrochloride Dose, and Angiography Results By Site

	Site No.		
	1	2	3
Angina class, %*			
None	26	17	16
I	3	1	4
II	26	20	21
III	34	40	44
IV	11	22	16
Sample size, No.	303	242	243
Daily propranolol dose, mg, %†			
20-40	34	16	15
60-80	35	31	31
120-240	25	46	49
>240	6	8	5
Sample size, No.	71	101	84
No. of major coronary vessels with ≥ 70% reduction in luminal diameter, %‡			
0	17	19	18
1	24	21	27
2	29	31	30
3	23	22	20
Left main	6	7	6
Sample size, No.	420	365	353

*This includes those patients who received coronary angiography for chronic stable angina, following a myocardial infarction, an episode of unstable angina, or coronary artery bypass surgery. $P < .01$ for differences among all sites, and for site 1 vs sites 2 and 3 combined.

†This includes only patients with angina classes I through IV receiving propranolol therapy. $P < .05$ for differences among sites; $P < .01$ for differences between site 1 vs sites 2 and 3 combined.

‡This excludes patients with no symptoms, nonspecific chest pain, congestive heart failure, and valvular disease. $P > .05$ for differences among all sites and for site 1 vs sites 2 and 3 combined.

These results can be viewed from several different perspectives. Physicians may argue that preservation of clinical independence and the full exercise of clinical judgment dictate that an indication be considered appropriate if a single physician believes it to be so. Payers might be less willing to provide reimbursement automatically for indications rated clearly inappropriate; they might, for example, see these as circumstances warranting second opinions. Before undergoing a procedure, a patient might want to know the relative appropriateness of the indication for it. Such information would allow the patient to add his or her preferences toward risk and benefit into the decision-making process. Some patients may prefer to undergo a procedure if the benefits only slightly outweigh the risks (eg, a 6 on a scale like the one we used); others may demand a wider margin.

We believe our findings on the differences among sites in the approaches physicians take toward coronary angiography can stimulate a productive educational dialogue. Data like these can help generate better agreement among physicians on when coronary angiography should and should not be performed. Our data on the level of appropriateness with which coronary angiography was performed in 1981 require confirmation for more recent years and younger populations and need

further validation by different methods. If confirmed, our finding that one of six coronary angiograms is performed inappropriately would suggest the need for an increased professional effort to improve the appropriateness with which this procedure is used.

This work was supported by the Commonwealth Fund, New York, the John A. Hartford Foundation, New York, the Health Care Financing Administration of the US Department of Health and Human Services, Washington, DC, the Pew Memorial Trust, Philadelphia, and The Robert Wood Johnson Foundation, Princeton, NJ.

This study could not have been completed without the support of the physicians and hospitals who permitted us to review their records. We would also like to thank the local affiliates and chapters of the American Heart Association in each of our sites for their support. Mary-Frances Flynn, MPH, Lois McCloskey, MPH, Joan Keesey, Barbara J. Genovese, Carole Fujihara, and Carole Oken, MA, deserve special thanks for their determination in seeing the fieldwork and data analysis through to their conclusions.

We are especially grateful to the physicians who served on the expert panel that wrestled with the difficult issues of appropriateness: Mortimer Buckley, MD, Ernie Chaney, MD, Gottlieb Friesinger, MD, J. Ward Kennedy, MD, John Kirklín, MD, Ceylon Lewis, MD, Joe Noble, MD, Harold Sox, Jr, MD, and William Wilson, MD.

References

- Gersh BJ, Kronmal RA, Frye RL, et al: Coronary arteriography and coronary artery bypass surgery: Morbidity and mortality in patients ages 65 years or older. *Circulation* 1983;67:483-490.
- Adams DF, Fraser DB, Abrams HL: The complications of coronary arteriography. *Circulation*

1973;48:609-618.

- Kennedy JW, and the Registry Committee of the Society for Cardiac Angiography: Complications associated with cardiac catheterization and angiography. *Cathet Cardiovasc Diagn* 1982;8:5-11.
- Kennedy JW, and the Registry Committee of the Society for Cardiac Angiography: Mortality related to cardiac catheterization and angiography. *Cathet Cardiovasc Diagn* 1982;8:323-340.
- Hansing CE: The risk and cost of coronary angiography: I. Cost of coronary angiography in Washington state. *JAMA* 1979;242:731-734.
- Chassin MR, Kosecoff J, Park RE, et al: *Indications for Selected Medical and Surgical Procedures: A Literature Review and Ratings of Appropriateness: Coronary Angiography*, publication R-32041-CWF-HF-HCFA-PMTR-RWJ. Santa Monica, Calif, The Rand Corp, 1986.
- Chassin MR, Brook RH, Park RE, et al: Variations in the use of medical and surgical services by the Medicare population. *N Engl J Med* 1986;314:285-289.
- Phibbs B: The abuse of coronary angiography. *N Engl J Med* 1979;301:1394-1396.
- Chassin MR, Kosecoff J, Park RE, et al: Does inappropriate use explain geographic variations in the use of health care services? A study of three procedures. *JAMA* 1987;258:2533-2537.
- Park RE, Fink A, Brook RH, et al: Physician ratings of appropriate indications for six medical and surgical procedures. *Am J Public Health* 1986;76:766-772.
- Park RE, Fink A, Brook RH, et al: *Physician Ratings of Appropriate Indications for Six Medical and Surgical Procedures*, publication R-3280-CWF-HF-PMTR-RWJ. Santa Monica, Calif, The Rand Corp, 1986.
- Kosecoff J, Chassin MR, Fink A, et al: Obtaining clinical data on the appropriateness of medical care in community practice. *JAMA* 1987;258:2538-2542.
- SAS User's Guide: Statistics, ed 5. Cary, NC, SAS Institute Inc, 1985.
- European Coronary Surgery Study Group: Prospective randomized study of coronary artery bypass surgery in stable angina pectoris. *Lancet* 1980;2:491-495.
- Takaro T, Hultgren HN, Lipton MJ, et al: The VA Cooperative Randomized Study of Surgery for Coronary Arterial Occlusive Disease: II. Subgroup with significant left main lesions. *Circulation* 1976;54:107-117.
- Murphy ML, Hultgren HN, Detke K, et al: Treatment of chronic stable angina: A preliminary report of survival data of the randomized Veterans Administration Cooperative Study. *N Engl J Med* 1977;297:621-627.
- CASS Principal Investigators and their associates: Coronary Artery Surgery Study (CASS): A randomized trial of coronary artery bypass surgery, survival data. *Circulation* 1983;68:939-950.
- Passamani E, Davis KB, Gillespie MJ, et al: A randomized trial of coronary artery bypass surgery: Survival of patients with a low ejection fraction. *N Engl J Med* 1985;312:1665-1671.
- Veterans Administration Coronary Artery Bypass Surgery Cooperative Study Group: 11-year survival in the Veterans Administration Randomized Trial of Coronary Bypass Surgery for Stable Angina. *N Engl J Med* 1984;311:1333-1339.
- Braunwald E: Effects of coronary-artery bypass grafting on survival: Implications of the Randomized Coronary-Artery Surgery Study. *N Engl J Med* 1983;309:1181-1184.
- Frishman WH, Furberg CD, Friedewald WT: β -Adrenergic blockade for survivors of acute myocardial infarction. *N Engl J Med* 1984;310:830-837.
- Lewis HD Jr, Davis JW, Archibald DG, et al: Protective effects of aspirin against acute myocardial infarction and death in men with unstable angina: Results of Veterans Administration Cooperative Study. *N Engl J Med* 1983;309:396-403.
- Smith B, Kennedy J: Thrombolysis in the treatment of acute transmural myocardial infarction. *Ann Intern Med* 1987;106:414-420.

