



# Developing Quality of Care Indicators for the Vulnerable Elderly

## The ACOVE Project

Far more people are surviving to old age than at any time in our history. Medical interventions can now prevent or postpone many of the health problems traditionally associated with old age. Furthermore, the goal of medical care for the elderly has progressed beyond survival to maximizing quality of life, yet little attention has been paid to the overall quality of medical care that older people receive. In fact, existing measures of quality or health status are often inappropriate for the elderly.

RAND Health has collaborated with Pfizer, Inc. to create the first quality-of-care assessment system for older persons. The goals of the Assessing Care of Vulnerable Elders (ACOVE) project were to define the population of vulnerable elderly, identify important medical conditions that affect this group, develop a set of evidence-based indicators of quality of care for this population, and design a tool to assess quality of care at the health system level.

### Previous Measures of Quality of Care in the Elderly Focused on Only a Few Conditions Appropriate to Elders

In the past, measures of quality of care in the elderly focused only on specific diseases or

**Key findings:**

- Little attention has been paid to the quality of health care that vulnerable elders and other older adults receive.
- RAND has developed a set of indicators to measure the quality of care received for 22 conditions that affect older adults.
- These indicators can be used to assess whether health plans are delivering care that meets minimum standards for quality.

aspects of care or targeted only a small proportion of the spectrum of older adults. Such focused approaches may not present a fair picture of overall quality, whereas broader systems of quality-of-care evaluation may exclude quality indicators for aspects of care that are most important to the well-being of older adults.

Quality of care may be more difficult to measure for older adults than for younger people. First, older adults show substantial variation in preferences for care. For example, older adults do not consistently prefer care that prolongs life, particularly if that care occurs at

**This Highlight summarizes RAND Health research reported in the following publications:**

*Annals of Internal Medicine*. 2001;135(Suppl.):641–758 is devoted to the ACOVE indicators. Articles cover the project overview, methods for developing the indicators, and the evidence supporting the quality indicators for 11 of the topics.

American College of Physicians–American Society of Internal Medicine web site, [www.acponline.org/sci-policy/acove](http://www.acponline.org/sci-policy/acove). Papers describe evidence supporting the other 11 topics.

- RAND RESEARCH AREAS
- CHILDREN AND ADOLESCENTS
- CIVIL JUSTICE
- EDUCATION
- ENERGY AND ENVIRONMENT
- HEALTH AND HEALTH CARE
- INTERNATIONAL AFFAIRS
- U.S. NATIONAL SECURITY
- POPULATION AND AGING
- PUBLIC SAFETY
- SCIENCE AND TECHNOLOGY
- SUBSTANCE ABUSE
- TERRORISM AND HOMELAND SECURITY
- TRANSPORTATION AND INFRASTRUCTURE

This product is part of the RAND Corporation research brief series. RAND research briefs present policy-oriented summaries of individual published, peer-reviewed documents or of a body of published work.

Corporate Headquarters  
1700 Main Street  
P.O. Box 2138  
Santa Monica, California  
90407-2138  
TEL 310.393.0411  
FAX 310.393.4818

© RAND 2004

the expense of comfort. Second, many ill older adults cannot advocate for themselves and may have no family members or friends to do so on their behalf. The ACOVE project endeavored to consider the heterogeneity and special needs of this population in designing a comprehensive set of quality assessment tools.

### Identifying Vulnerable Elders

We defined vulnerable elders as those persons 65 years of age and older who are at high risk for death or functional decline and devised a set of criteria to identify members of that group for measurement. We excluded the use of utilization data as selection criteria because selecting individuals solely on the basis of the health care they have received might miss an important component of the population—those who are undertreated or underdiagnosed. In addition, based on a longitudinal analysis of the Medicare Current Beneficiary Survey, we determined that self-rated functional status was a more important predictor of functional decline and death than are specific clinical conditions. We designed a brief telephone survey that assessed age, self-rated health, limitations in physical capability, and functional limitations. Individuals who were identified as vulnerable on the survey were more than four times as likely to die or experience functional decline over a two-year period as those who were not so identified. According to these criteria, 32 percent of a nationally representative sample of elders was defined as vulnerable.

### Developing and Implementing the Quality-of-Care Indicators

To develop the system, a national panel of geriatrics experts identified the medical conditions prevalent among older adults that contribute most to morbidity, mortality, and functional decline; that could be measured; and for which effective methods of treatment or prevention are available. Using these criteria, our advisory committee chose 22 topics, including diseases, syndromes, physiological impairments, and clinical situations, for which quality-of-care indicators could be developed for this population (see box, “ACOVE Topics”). According to national surveys, the prevalence of the selected conditions ranged from 10 percent to 50 percent among community-dwelling older adults.

For each condition, the RAND team developed quality-of-care indicators. Quality of care can be measured using either processes of care or outcomes of care. We chose to use process measures because processes are a more efficient measure of quality of care and are amenable to direct change. However, associations between care processes and outcomes have been supported by high-quality research in only a few

### ACOVE Topics

Appropriate Use of Medication  
Chronic Pain  
Continuity and Coordination of Care  
Dementia  
Depression  
Diabetes Mellitus  
End-of-Life Care  
Falls and Mobility Problems  
Hearing Loss  
Heart Failure  
Hospital Care  
Hypertension  
Ischemic Heart Disease  
Malnutrition  
Osteoarthritis  
Osteoporosis  
Pneumonia  
Pressure Ulcers  
Preventive Care  
Stroke and Atrial Fibrillation  
Urinary Incontinence  
Visual Impairment

instances, and those studies often exclude vulnerable elders. As we developed our quality indicators, our clinical experts assessed the available evidence for its applicability to vulnerable elders.

A set of potential quality indicators was developed for each of the 22 topics using existing guidelines and expert opinion. These indicators covered four domains of care:

- Prevention
- Diagnosis
- Treatment
- Follow-up.

Our team performed structured literature reviews to assess the evidence supporting a link between each of the proposed care processes and improved outcomes in older

### **What Does a Quality Indicator Look Like?**

#### **Quality Indicator 1 for Dementia: Cognitive and Functional Screening**

IF a vulnerable elder is admitted to a hospital or is new to a physician practice, THEN multidimensional assessment of cognitive ability and assessment of functional status should be documented BECAUSE screening for dementia can lead to early detection and initiation of treatment that may delay further progression.

adults. The proposed quality indicators and the supporting literature were reviewed by independent panels of experts in geriatric care and the medical specialties, who assessed their validity and feasibility using a variation of the RAND/UCLA Appropriateness Method for developing guidelines to measure the appropriateness of medical care. Based on the panels' ratings, we developed a set of 236 quality indicators covering the 22 topics (see box, "What Does a Quality Indicator Look Like?"). The quality indicators accepted by the panels, as well as the supporting literature reviews, were further evaluated by the American College of Physicians–American Society of Internal Medicine Aging Task Force before acceptance.

The ACOVE quality indicators reflect the concerns of the geriatric patient population they are intended to serve. Many of the indicators focus on the transfer of information between the provider and the patient or the patient's proxy; others focus on detecting and treating conditions that are underdetected in the elderly. These conditions include dementia, depression, and functional impairments. Furthermore, underlying the indicators are considerations such as advance care planning and informed consent to treatment.

To implement the quality indicators in care settings, we developed instruments to abstract medical records, interview patients or proxies, and evaluate administrative data. Pilot testing of most of the indicators was completed in two managed care plans. In recognition of the unique concerns of the vulnerable elders, we considered several factors that might mitigate the decision to apply a particular indicator to the care of a particular patient. These factors included patient preferences to avoid hospitalization or surgery and conditions such as advanced dementia or a documented

terminal prognosis (which might lead providers to withhold some otherwise indicated elements of care). In addition, when implementing the indicators, evaluators considered local guidelines and resources available for care.

### **When the Indicators Can and Cannot Be Used**

As we explained above, the ACOVE indicator system is the first quality assessment system specifically designed for the care of ill older adults. However, we must note two important features about the ACOVE indicators.

First, the indicators are not practice guidelines. Practice guidelines aim to define optimal or ideal care in the context of complex decisionmaking. In contrast, quality indicators set a minimal standard for acceptable care—standards that, if not met, almost ensure that the care is of poor quality.

Second, the ACOVE indicators are designed to evaluate health care at the system (or plan) level, not at the individual level. This means that they cannot be used to evaluate individual physicians. Furthermore, the indicators have not yet been tested with sufficient numbers of patients to ensure their utility for evaluating the care of an individual patient or the treatment of a single condition. Rather, the indicators currently allow us to assess the overall care delivered to vulnerable elders by their health care plans or medical groups and thus could be used to identify areas in need of improvement.

### **What Next?**

RAND researchers have now implemented the ACOVE indicators to assess the quality of care received by vulnerable elders who live at home and are enrolled in one of two managed care organizations throughout the United States. The researchers assessed the quality of care for both geriatric and general medical conditions. Also assessed was the quality of pharmacological care—prescribing medications—for vulnerable elders. A follow-up research highlight, "The Quality of Care Older Adults Receive," reviews their findings.

Currently, ACOVE is in its second phase. ACOVE-2 aims to improve the medical care physicians provide to older adults by developing and testing interventions to increase performance on selected quality indicators. These indicators correspond to geriatric care processes that were the least well carried out in the quality assessment phase. The goal of ACOVE-2 is to develop methods for changing clinical practice that will be tailored to specific practice settings. ■

---

Abstracts of all RAND Health publications and full text of many research documents can be found on the RAND Health web site at [www.rand.org/health](http://www.rand.org/health). The RAND Corporation is a nonprofit research organization providing objective analysis and effective solutions that address the challenges facing the public and private sectors around the world. RAND's publications do not necessarily reflect the opinions of its research clients and sponsors. RAND® is a registered trademark.

**RAND Offices** Santa Monica • Washington • Pittsburgh • New York • Doha • Berlin • Cambridge • Leiden



# HEALTH

CHILD POLICY  
CIVIL JUSTICE  
EDUCATION  
ENERGY AND ENVIRONMENT  
HEALTH AND HEALTH CARE  
INTERNATIONAL AFFAIRS  
NATIONAL SECURITY  
POPULATION AND AGING  
PUBLIC SAFETY  
SCIENCE AND TECHNOLOGY  
SUBSTANCE ABUSE  
TERRORISM AND  
HOMELAND SECURITY  
TRANSPORTATION AND  
INFRASTRUCTURE

This PDF document was made available from [www.rand.org](http://www.rand.org) as a public service of the RAND Corporation.

This product is part of the RAND Corporation research brief series. RAND research briefs present policy-oriented summaries of individual published, peer-reviewed documents or of a body of published work.

The RAND Corporation is a nonprofit research organization providing objective analysis and effective solutions that address the challenges facing the public and private sectors around the world.

---

## Support RAND

[Browse Books & Publications](#)

[Make a charitable contribution](#)

## For More Information

Visit RAND at [www.rand.org](http://www.rand.org)

Explore [RAND Health](#)

View [document details](#)

## Limited Electronic Distribution Rights

This document and trademark(s) contained herein are protected by law as indicated in a notice appearing later in this work. This electronic representation of RAND intellectual property is provided for non-commercial use only. Permission is required from RAND to reproduce, or reuse in another form, any of our research documents for commercial use.