

U. S. AIR FORCE  
PROJECT RAND  
RESEARCH MEMORANDUM

SOME NOTES FOR SIMPLE PAVLOVIAN LEARNING

A. S. Householder

RM-678

10 September 1951

Assigned to \_\_\_\_\_

This is a working paper. It may be expanded,  
modified, or withdrawn at any time.



**SUMMARY:** Some neural nets capable of mediating simple Pavlovian reflexes in their qualitative features are represented, along with some suggestions toward further elaboration.

SOME NOTES FOR SIMPLE PAVLOVIAN LEARNING

by A. S. Householder

In the accompanying figures are exhibited some nets representing some of the qualitative features of the simplest types of Pavlovian reflex learning. It is perhaps clear how, by inserting internuncials and adding circuits, the quantitative characteristics can be adjusted. Admittedly a great many such additions might be required before achieving any degree of realism.

Fig. 1 represents the simplest form of conditioning with extinction. Here and hereafter the threshold is 1 or irrelevant unless inserted; and S and S' denote the unconditioned and conditioned stimuli, respectively. In this net the occurrence of S'(t) S(t + 1) activates the cycle, after which S'(T) implies R(T + 2). However any occurrence of S'(t) ~ S(t + 1) inactivates the cycle. Thus a single trial establishes conditioning, but a single failure to reinforce will extinguish it.

Fig. 2 shows one step in the elaboration. Here one trial activates C<sub>1</sub> but does not yet set up the conditioning; a second trial is required, at which time C<sub>2</sub> is activated and S' (via an internuncial for timing purposes) is made liminal. A single nonreinforcement will inactivate C<sub>2</sub>; a second will inactivate also C<sub>1</sub>.

If C<sub>1</sub>, instead of C<sub>2</sub>, were connected to R, then a single trial would establish conditioning, but unstably, in the sense that a single nonreinforcement

would extinguish it. Two consecutive trials would increase the stability, necessitating two consecutive nonreinforcements to extinguish it. This is readily generalized by taking 2 cycles in stages, of which the  $m$ -th is connected to R. Then conditioning would be established after  $m$  trials, but each further trial increases the stability, up to the number  $n - m$  of additional stages.

Fig. 3 represents the first aspect of the delayed reflex, with one-stage conditioning. Note that  $S(t)$  always implies  $R(t + 1)$ . Now the occurrence of  $S'(t) S(t + 1)$  activates  $C_1$  as in Fig. 1, so that subsequent occurrence of  $S'(\tau)$  implies  $R(\tau + 2)$ . However if  $\sim S(\tau + 1)$ , then the cycle is inactivated and the conditioning extinguished. Also the occurrence of  $S'(t) S(t + 2)$  activates  $C_2$  so that subsequent occurrence of  $S'(\tau)$  implies  $R(\tau + 3)$ .

The most interesting feature of the delayed reflex, however, in Pavlov's description is this. Continued reinforcement of the delayed reflex (corresponding to activation of  $C_2$  and subsequent stages) has the effect that the time of occurrence of R moves progressively forward; non-reinforcement first increases the delay before continued nonreinforcement finally extinguishes the conditioning.

To achieve this in its most elementary form, add another stage to  $C_2$  as in Fig. 2, but connect this cycle to  $I_2$  with an inhibitor and to  $I_1$  with an excitor. Then  $S'(t)$  is sufficient to produce  $R(t + 2)$  if either this new cycle is active, or if  $C_1$  is active, whereas if  $C_2$  alone is active, then  $S'(t)$  will produce  $R(t + 3)$ .

Fig. 4 gives the phenomenon of generalization. It is supposed that  $S_1'$  and  $S_2'$  are affected by different degrees of intensity, or different qualities of the same type of stimulus (e.g., distinct tones). Either conjunction  $S_1'(t) S(t + 1)$  or  $S_2'(t) S(t + 1)$  is sufficient to activate both cycles  $C_1$  and  $C_2$ , after which the occurrence of either  $S_1'(\tau)$  or of

$S_2'(\tau)$  is sufficient for  $R(\tau + 2)$ . However, the non-reinforced presentation of  $S_1'$  above, that is to say  $S_1'(\tau) \sim S(\tau + 1)$ , inactivates  $C_1$ , but not  $C_2$ .

A more realistic representation requires obviously more than two degrees or qualities, but should have also the following features: Suppose the afferents are  $S_1', \dots, S_2'$ , in the order of degrees of intensity or of a simple ordering of the qualities. Progressive reinforcements of a particular  $S_\alpha'$  should establish  $S_\alpha'$  sooner than other  $S_\beta'$  and for establishing  $S_\beta'$  by reinforcing  $S_\alpha'$ , the required number of reinforcements should increase with  $(\beta - \alpha)$ . Conversely, in the subsequent extinction of a given  $S_\beta'$ , more non-reinforcements should be required for small  $(\beta - \alpha)$  than for large. Again, as extinction proceeds by non-reinforcement of  $S_\beta'$ , extinction of any  $S_\gamma'$  should proceed at about the same relative rate when  $\beta$  lies between  $\gamma$  and  $\alpha$ ; whereas when  $\gamma$  lies between  $\beta$  and  $\alpha$  the rate at which  $S_\gamma'$  is extinguished should depend upon the relative values of  $(\beta - \gamma)$  and  $(\gamma - \alpha)$ .






Fig. 2— Two-stage conditioning




Fig. 3 — Simple delayed reflex






Fig. 4—Generalization



