



CHILDREN AND FAMILIES
EDUCATION AND THE ARTS
ENERGY AND ENVIRONMENT
HEALTH AND HEALTH CARE
INFRASTRUCTURE AND
TRANSPORTATION
INTERNATIONAL AFFAIRS
LAW AND BUSINESS
NATIONAL SECURITY
POPULATION AND AGING
PUBLIC SAFETY
SCIENCE AND TECHNOLOGY
TERRORISM AND
HOMELAND SECURITY

The RAND Corporation is a nonprofit institution that helps improve policy and decisionmaking through research and analysis.

This electronic document was made available from www.rand.org as a public service of the RAND Corporation.

Skip all front matter: [Jump to Page 1](#) ▼

Support RAND

[Browse Reports & Bookstore](#)

[Make a charitable contribution](#)

For More Information

Visit RAND at www.rand.org

Explore the [RAND Corporation](#)

View [document details](#)

Limited Electronic Distribution Rights

This document and trademark(s) contained herein are protected by law as indicated in a notice appearing later in this work. This electronic representation of RAND intellectual property is provided for non-commercial use only. Unauthorized posting of RAND electronic documents to a non-RAND website is prohibited. RAND electronic documents are protected under copyright law. Permission is required from RAND to reproduce, or reuse in another form, any of our research documents for commercial use. For information on reprint and linking permissions, please see [RAND Permissions](#).

This report is part of the RAND Corporation research report series. RAND reports present research findings and objective analysis that address the challenges facing the public and private sectors. All RAND reports undergo rigorous peer review to ensure high standards for research quality and objectivity.

Maternal Depression: Implications for Systems Serving Mother and Child

Lisa Sontag-Padilla, Dana Schultz, Kerry A. Reynolds, Susan L. Lovejoy, & Ray Firth

Implications

- Identification of and treatment for maternal depression may improve short- and long-term outcomes for mothers and their children.
- Untreated maternal depression has been associated with negative outcomes in the areas of employment and income, which has potential cost implications for the public assistance system.
- Untreated maternal depression has been associated with poor health outcomes for both mothers and children, which may be associated short-term and long-term costs to the health system.
- Maternal depression is also linked to early childhood developmental delays, which put children at greater risk of needing early intervention services.
- Children of depressed mothers are more at risk for early cognitive developmental delays and poor academic performance, which could increase needs for special education services.
- Untreated maternal depression may increase risk for child maltreatment and neglect, which may increase need for subsequent involvement of the child welfare system.
- Ultimately, reducing the prevalence of maternal depression may have short- and long-term financial implications for the publicly funded systems that serve depressed mothers and their children.

SUMMARY ■

This report highlights evidence on the impact of maternal depression on the mother and child as it relates to the public-sector systems that serve them and discusses potential short- and long-term cost implications. The goal of this brief is to serve as a source of information for state and local policymakers and practitioners concerned with child and family outcomes, and to inform them of the evidence connecting maternal depression and negative outcomes for mother and child. The overview of the evidence is not meant to be exhaustive; rather, this brief highlights evidence most relevant to the impact of untreated maternal depression on both the mother and the child and the potential cost implications for systems that serve those families affected by depression. Due to the lack of clear data implicating the specific costs of maternal depression to each public-sector system, this brief focuses on the conceptual links between maternal depression and negative outcomes, as well as the hypothesized impact on each system.

BACKGROUND


Depression affects millions of Americans each year and bears significant societal and financial costs (Kessler, 2012; Wang et al., 2006). However, it is estimated that only 25 percent of individuals with depression receive appropriate care (Young et al., 2001). Given that women are almost twice as likely to experience depression compared to men (Gaynes et al., 2005;

Kessler et al., 2003; Kessler et al., 2005) and that the majority of women age 15 to 50 have children (U.S. Census Bureau, 2012), maternal depression is an important and potentially costly issue.

Each year, more than 400,000 babies are born to women experiencing depression (Earls, 2010) and an estimated 15 million children live with a mother experiencing depression (National Research Council and Institute of Medicine, 2009). Untreated maternal depression has potentially serious consequences for a woman's overall well-being, her functioning as a mother, the family's functioning, and her child's development (Cummings et al., 2008; Elgar et al., 2007; Goodman & Gotlib, 1999; Lim, Wood, & Miller, 2008; Onunaku, 2005; Field, 2000). However, identification of and treatment for maternal depression may improve short- and long-term outcomes for both mothers and their children (Center on the Developing Child at Harvard University, 2009; Cicchetti, Rogosch, & Toth, 2000; Clark, Tluczek, & Wenzel, 2003).

In turn, reducing the prevalence of maternal depression may also have short- and long-term financial implications for the publicly funded systems that serve depressed mothers and their children. Evidence on the cost-benefit ratio of treating maternal depression is limited to local estimates that include only some of the potential costs discussed in this brief (Diaz & Chase, 2010; Lynch & Harrington, 2003). However, these estimates suggest that the public benefits of treating a case of maternal depression are likely to exceed the costs. In an effort to provide context to the potential impact of untreated maternal depression, this report summarizes evidence on the two-

Public Sector Systems Affected by Maternal Depression



RAND RR404-1

generational (mother and child) negative implications of maternal depression on different public-sector systems that serve mothers and children (see Figure). While this overview is not meant to be exhaustive, we highlight evidence relevant to the impact of untreated maternal depression on both the mother and the child and describe the potential short- and long-term cost implications for some of the systems that may be affected.

IMPLICATIONS OF MATERNAL DEPRESSION ON PUBLIC-SECTOR SYSTEMS

In this section, we summarize some of the evidence on the impact of maternal depression (on the mother and child) as it relates to different public-sector systems, including public assistance, physical health, early intervention, education, and child welfare. In each section, we also highlight the potential short- and long-term impact on the public-sector system.

Public Assistance System

Untreated maternal depression has been associated with negative outcomes in the areas of employment and income, which has potential cost implications for the public assistance system. Depressed mothers are more likely to be unemployed (20 percent vs. 8 percent) and less likely to be employed full time (40 percent vs. 52 percent) when compared to nondepressed mothers (Ertel, Rich-Edwards, & Koenen, 2011). However, because these data were cross-sectional, the correlation remains unclear; i.e., the extent to which depression causes unemployment versus unemployment leading to depression. Depressed mothers may also have difficulty getting and keeping a job, leading to lower income and a greater need for public assistance (Ertel, Rich-Edwards, & Koenen, 2011; Kawakami et al., 2012; Lepine & Briley, 2011; Turney, 2012). For the general population, depression has been shown to be predictive of greater work disability in the short term and of lower income over time (Kawakami et al., 2012; Lepine & Briley, 2011).

Moreover, treatment of depression has been shown to improve work productivity and decrease absenteeism (Rost, Smith, & Dickinson, 2004; Schoenbaum et al., 2002; Wang et al., 2006). Given the costs associated with unemployment and low income—such as Temporary Assistance to Needy Families; unemployment benefits; food stamps; assistance from Women, Infants and Children—reducing the occurrence of maternal

Reducing the prevalence of maternal depression may have short- and long-term financial implications for the publicly funded systems that serve depressed mothers and their children.

depression has potential cost savings implications for the public assistance system.

Physical Health Care System

Untreated maternal depression has been associated with poor health outcomes for both mothers and children. For the mother, depression may increase the likelihood of certain disease conditions both during the maternal part of her lifecycle as well as throughout her entire life course. Depressed women are more likely to have cardiovascular disease (Farr et al., 2011; Pan et al., 2011; Wuslin, 2004), stroke (Rexrode, 2010) and type-2 diabetes (Bowers et al., 2013; Farr et al., 2011; Mezuk et al., 2008; Pan et al., 2011). Some studies found women with major depression to be 3.6 times more likely than nondepressed women to have diabetes and chronic disease risk factors (Farr et al., 2011) and 1.45 times more likely to have a stroke (Pan et al., 2011).

Maternal depression also may lead indirectly to poor health outcomes for the child, such as increased risk for infant hospitalization (Chung et al., 2004) and asthma morbidity (Pak & Jackson Allen, 2012). Babies born to depressed women are also more likely to be premature (Li, Liu, & Odouli, 2009; Orr, James, & Blackmore Prince, 2002) and are at greater risk of being small for gestational age (U.S. Department of Health and Human Services, Office of Women's Health, 2009); both of these factors increase the likelihood of neonatal intensive care unit (NICU) stays, which are costly. One analysis estimated that the cost of a night in the NICU was nearly 2.4 times the cost of a regular nursery night (Adams et al., 2002) with NICU stays typically lasting several weeks for most babies. Identifying and treating maternal depression provides one means of reducing prevalence of certain disease conditions and poor birth

outcomes to potentially offset some of the associated short-term and long-term costs to the health system.

Early Intervention System

Maternal depression is also linked to early childhood developmental delays. Depressed mothers may be more likely to have children with early childhood developmental delays and thus require services through the early intervention system. Research indicates that maternal depression poses a serious risk to the quality of the parent-child relationship (Cummings et al., 2008; Elgar et al., 2007; Goodman & Gotlib, 1999; Lim et al., 2008), which in turn can threaten a child's physical, social, and cognitive development during early childhood (Davies, Winter, & Cicchetti, 2006; Sroufe et al., 2005). For instance, infants and young children of depressed mothers have increased risk for social and emotional problems (Moore, Cohn, & Campbell, 2001; Whitaker, Orzol, & Kahn, 2006) and delays or impairments in cognitive and linguistic development (Grace, Evindar, & Stewart, 2003; Downey & Coyne, 1990). These impairments or delays put children at greater risk of needing early intervention services related to cognitive and language delays. Additionally, mothers of children with developmental delays are more likely to experience depression (National Research Council and Institute of Medicine, 2009), potentially perpetuating the negative cycle and increasing short- and long-term costs associated with early intervention services.

Education System

Untreated maternal depression can potentially have negative effects on child-level education outcomes as well. Children of depressed mothers may be less likely to enter school ready to learn because of increased risk for early cognitive developmen-

tal delays and poor academic performance (Kersten-Alvarez et al., 2012; Sohr-Preston & Scaramella, 2006). This, in turn, may increase children's needs for special education services. For instance, children of depressed mothers show lower IQs (significant difference of 4.5 points) during the early childhood years (Evans et al., 2012), potentially leading to greater need for special education services. Given that quality special education costs on average 1.3 times as much as teaching students without special needs (Augenblick and Associates, Inc., 2009), the increased risk for cognitive developmental delays and poor academic performance due to maternal depression may incur downstream costs to the public education system.

Child Welfare System

Untreated maternal depression may increase risk for child maltreatment and neglect (Collishaw et al., 2007; Hazen et al., 2006; Koverola et al., 2005; Taylor et al., 2009) due to the lack of attentiveness and responsiveness associated with depression (Diego, Field, & Hernandez-Reif, 2005). This, in turn, may increase risk for subsequent involvement of the child welfare system. Given the high costs associated with child victims of maltreatment—e.g., estimated lifetime cost is approximately \$210,012 per child (Fang et al., 2012)—reducing some of the risk for child maltreatment through prevention or treatment of maternal depression may help offset costs within the child welfare system.

CONCLUSIONS

As outlined here, untreated maternal depression may have far-reaching negative implications for mothers and children, many of which may be associated with high costs for multiple public-sector service systems. Despite the frequency of depression among new mothers, large numbers of affected women and their children go unidentified, and an estimated 85 percent do not seek treatment (Center on the Developing Child at Harvard University, 2009). Although the potential costs associated with untreated maternal depression may be reduced or eliminated by focusing additional resources on the identification and treatment of depression, prevention efforts to reduce risk for, and incidence of, maternal depression may prove to be just as valuable if not more cost-effective (Center on the Developing Child at Harvard University, 2009). However, due to a dearth of data on the estimated costs of maternal depression to the different public-sector service systems, our conclusions are based on the conceptual evidence linking maternal depression with various negative outcomes. That is, while it is established that maternal depression has multiple and far-reaching negative effects on the mother's work productivity, employment, income, and physical health, along with the child's physical, cognitive, and behavioral development and the parent-child relationship, the magnitude of the direct impact on the associated public-sector service systems remains unknown. Despite these limitations, the evidence suggests that identifying depressed mothers early and reaching them with effective, evidence-based treatments may offset some short- and long-term costs for various public-sector systems by reducing the likelihood that mothers and children will need service or supports.

About This Report

This brief was developed as a supplemental component of the *Helping Families Raise Healthy Children* initiative implemented in Allegheny County, Pennsylvania, which successfully implemented depression screening in early intervention, developed cross-system referral processes, engaged caregivers in services, and increased local capacity for providing relationship-based services by addressing many of the barriers identified in earlier stages of the collaborative's work (Schultz et al., 2012; 2013).

The *Helping Families Raise Healthy Children* initiative and the preparation of this brief were supported by the Robert Wood Johnson Foundation Local Funding Partnership grant program, the Highmark Foundation as the nominating funder, and other local funding partners including UPMC Health Plan, The Pittsburgh Foundation, The Fine Foundation, FISA Foundation, and the Jewish Healthcare Foundation, with additional support from the Allegheny County Department of Human Services, Office of Behavioral Health, and the Pennsylvania Department of Public Welfare. The research was conducted within RAND Health, a division of the RAND Corporation. The RAND Health Quality Assurance process employs peer reviewers, including at least one reviewer who is external to the RAND Corporation. A profile of RAND Health, abstracts of its publications, and ordering information can be found at www.rand.org/health.

References

- Adams, E. K., Miller, V. P., Ernst, C., Nishimura, B. K., Melvin, C., & Merritt, R. (2002). Neonatal health care costs related to smoking during pregnancy. *Health Economics*, *11*(3), 193–206.
- Augenblick, P. and Associates, Inc. (2009). *Costing out the resources needed to meet Pennsylvania's public education goals*.
- Bowers, K., Laughon, S. K., Kim, S., Mumford, S. L., Brite, J., Kiely, M., & Zhang, C. (2013). The association between a medical history of depression and gestational diabetes in a large multi-ethnic cohort in the United States. *Paediatric and Perinatal Epidemiology*, *27*(4), 323–328.
- Center on the Developing Child at Harvard University (2009). *Maternal Depression Can Undermine the Development of Young Children*. Working Paper No. 8.
- Chung, E. K., McCollum, K. F., Elo, I. T., Lee, H. J., & Culhane, J. F. (2004). Maternal depressive symptoms and infant health practices among low-income women. *Pediatrics*, *113*(6), e523–e529.
- Cicchetti, D., Rogosch, F. A., & Toth, S. L. (2000). The efficacy of toddler-parent psychotherapy for fostering cognitive development in offspring of depressed mothers. *Journal of Abnormal Child Psychology*, *28*, 135–148.
- Clark, R., Tluczek, A., & Wenzel, A. (2003). Psychotherapy for postpartum depression: A preliminary report. *American Journal of Orthopsychiatry*, *73*, 441–454.
- Collishaw, S., Dunn, J., O'Connor, T. G., Golding, J., & the Avon Longitudinal Study of Parents and Children Study. (2007). Maternal childhood abuse and offspring adjustment over time. *Development and Psychopathology*, *9*, 367–383.
- Cummings, E. M., Schermerhorn, A. C., Keller, P. S., & Davies, P. T. (2008). Parental depressive symptoms, children's representations of family relationships, and child adjustment. *Social Development*, *17*, 278–305.
- Davies, P. T., Winter, M. A., & Cicchetti, D. (2006). The implications of emotional security theory for understanding and treating childhood psychopathology. *Development and Psychopathology*, *18*, 707–735.
- Diaz, J. Y., & Chase, R. (2010). *The cost of untreated maternal depression*. St. Paul, MN: Wilder Research.
- Diego, M. A., Field, T., & Hernandez-Reif, M. (2005). Prepartum, postpartum and chronic depression effects on neonatal behavior. *Infant Behavioral Development*, *28*(5), 155–164.
- Downey, G., & Coyne, J. C. (1990). Children of depressed parents: An integrative review. *Psychological Bulletin*, *108*, 50–76.
- Earls, M. F. (2010). Incorporating recognition and management of perinatal and postpartum depression into pediatric practice. *Pediatrics*, *126*(5), 1032–1039.
- Elgar, F. J., Mills, R. S. L., McGrath, P. J., Waschbusch, D. A., & Brownridge, D. A. (2007). Maternal and paternal depressive symptoms and child maladjustment: The mediating role of parental behavior. *Journal of Abnormal Child Psychology*, *35*, 943–955.
- Ertel, K. A., Rich-Edwards, J. W., & Koenen, K. C. (2011). Maternal depression in the United States: Nationally representative rates and risks. *Journal of Women's Health*, *20*(11), 1609–1617.
- Evans, J., Melotti, R., Heron, J., Ramchandani, P., Wiles, N., Murray, L., & Stein, A. (2012). The timing of maternal depressive symptoms and child cognitive development: A longitudinal study. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, *53*(6), 632–640.
- Fang, X., Brown, D. S., Florence, C. S., & Mercy, J. A. (2012). The economic burden of child maltreatment in the United States and implications for prevention. *Child Abuse & Neglect*, *36*, 156–165.
- Farr, S. L., Hayes, D. K., Bitsko, R. H., Bansil, P., & Dietz, P. M. (2011). Depression, diabetes, and chronic disease risk factors among US women of reproductive age. *Preventing Chronic Disease*, *8*(6), A119. As of December 2, 2011: http://www.cdc.gov/pcd/issues/2011/nov/10_0269.htm
- Field, T. M. (2000). Infants of depressed mothers. In Johnson, S. L., *Stress, coping, and depression*, 3–23. Mahwah, NJ: L. Erlbaum Associates.
- Gaynes, B. N., Gavin, N., Meltzer-Brody, S., Lohr, K. N., Swinson, T., Gartlehner, G., et al. (2005). *Perinatal depression: Prevalence, screening accuracy, and screening outcomes*. Evidence report/technology assessment No. 119. AHRQ Publication No. 05-E006–2. Rockville, MD: Agency for Healthcare Research and Quality.

- Goodman, S. H., & Gotlib, I. H. (1999). Risk for psychopathology in the children of depressed mothers: A developmental model for understanding mechanisms of transmission. *Psychological Review*, *106*(3), 458–490.
- Grace, S. L., Evindar, A., & Stewart, D. E. (2003). The effect of postpartum depression on child cognitive development and behavior: A review and critical analysis of the literature. *Archives of Women's Mental Health*, *6*(4), 263–274.
- Hazen, A. L., Connelly, C. D., Kelleher, K. J., Barth, R. P., & Landsverk, J. A. (2006). Female caregivers' experiences with intimate partner violence and behavior problems in children investigated as victims of maltreatment. *Pediatrics*, *117*, 99–109.
- Kawakami, N., Abdulghani, E. A., Alonso, J., Bromet, E. J., Bruffaerts, R., Caldas-de-Almeida, J. M., et al. (2012). Early-life mental disorders and adult household income in the world mental health surveys. *Biological Psychiatry*, *72*(3), 228–237.
- Kersten-Alvarez, L. E., Hosman, C. M., Riksen-Walraven, J. M., van Doesum, K. T., Smeekens, S., & Hoefnagels, C. (2012). Early school outcomes for children of postpartum depressed mothers: Comparison with a community sample. *Child Psychiatry and Human Development*, *43*(2), 201–218.
- Kessler, R. C. (2012). The costs of depression. *The Psychiatric Clinics of North America*, *35*(1), 1–14.
- Kessler, R. C., Berglund, P., Demler, O., Jin, R., Koretz, D., Merikangas, K. R., et al. (2003). The epidemiology of major depressive disorder: Results from the National Comorbidity Survey Replication (NCS-R). *Journal of the American Medical Association* *289*, 3095–3105.
- Kessler, R. C., Chiu, W. T., Demler, O., & Walters, E. E. (2005). Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, *62*(6), 617–627.
- Koverola, C., Papas, M. A., Pitts, S., Murtaugh, C., Black, M. M., & Dubowitz, H. (2005). Longitudinal investigation of the relationship among maternal victimization, depressive symptoms, social support, and children's behavior and development. *Journal of Interpersonal Violence*, *20*, 1523–1546.
- Lepine, J. P., & Briley, M. (2011). The increasing burden of depression. *Neuropsychiatric Disease and Treatment*, *7*(Suppl 1), 3–7.
- Li, D., Liu, L., & Odouli, R. (2009). Presence of depressive symptoms during early pregnancy and the risk of preterm delivery: A prospective cohort study. *Human Reproduction*, *24*(1), 146–153.
- Lim, J., Wood, B. L., & Miller, B. D. (2008). Maternal depression and parenting in relation to child internalizing symptoms and asthma disease activity. *Journal of Family Psychology*, *22*, 264–273.
- Lynch, T., & Harrington, J. (2003). *Benefit cost analysis of the maternal depression project in Gadsden County, Florida*. Tallahassee, FL: Center for Economic Forecasting and Analysis (CEFA).
- Mezuk, B., Eaton, W. W., Albrecht, S., & Golden, S. H. (2008). Depression and type-2 diabetes over the lifespan: A meta-analysis. *Diabetes Care* *31*, 2383–2390.
- Moore, G. A., Cohn, J. F., & Campbell, S. B. (2001). Infant affective responses to mother's still face at 6 months differentially predict externalizing and internalizing behaviors at 18 months. *Developmental Psychology*, *37*(5), 706–714.
- National Research Council and Institute of Medicine. (2009). *Depression in parents, parenting, and children: Opportunities to improve identification, treatment, and prevention*, Committee on Depression, Parenting Practices, and the Healthy Development of Children. Board on Children, Youth, and Families. Division of Behavioral and Social Sciences and Education. Washington, DC: The National Academies Press.
- Onunaku, N. (2005). *Improving maternal and infant mental health: Focus on maternal depression*. Los Angeles, CA: National Center for Infant and Early Childhood Health Policy at UCLA.
- Orr, S. T., James, S. A., & Blackmore Prince, C. (2002). Maternal prenatal depressive symptoms and spontaneous preterm births among African-American women in Baltimore, Maryland. *American Journal of Epidemiology*, *156*, 797–802.
- Pak, L., & Allen, P. J. (2012). The impact of maternal depression on children with asthma. *Pediatric Nursing*, *38*(1), 11–19, 30.
- Pan, A., Sun, Q., Okereke, O. I., Rexrode, K. M., & Hu, F. B. (2011). Depression and risk of stroke morbidity and mortality: A meta-analysis and systematic review. *Journal of the American Medical Association*, *306*, 1241–1249.
- Rexrode, K. M. (2010). Emerging risk factors in women. *Stroke: A Journal of Cerebral Circulation*, *41*(10 Suppl), S9–S11.

- Rost, K., Smith, J. L., & Dickinson, M. (2004). The effect of improving primary care depression management on employee absenteeism and productivity: A randomized trial. *Medical Care*, *42*, 1202–1210. [PubMed: 15550800]
- Schoenbaum, M., Unutzer, J., McCaffrey, D., Duan, N., Sherbourne, C., & Wells, K. B. (2002). The effects of primary care depression treatment on patients' clinical status and employment. *Health Services Research*, *37*(5), 1145–1158.
- Schultz, D., Reynolds, K., Sontag-Padilla, L., Lovejoy, S. L., Firth, R., Schake, P., et al. *A toolkit for implementing parental depression screening, referral, and treatment across systems*, Santa Monica, CA: RAND Corporation, TL-102-CCBHO. As of July 2, 2013: <http://www.rand.org/pubs/tools/TL102>
- Schultz, D., Reynolds, K., Sontag-Padilla, L., Lovejoy, S. L., Firth, R., & Pincus, H. A. (2013). *Transforming systems for parental depression and early childhood developmental delays: Findings and lessons learned from the Helping Families Raise Healthy Children Initiative*. Santa Monica, CA: RAND Corporation, RR-122-CCBHO. As of July 24, 2013: http://www.rand.org/pubs/research_reports/RR122
- Sohr-Preston, S. L., & Scaramella, L. V. (2006). Implications of timing of maternal depressive symptoms for early cognitive and language development. *Clinical Child and Family Psychology Review*, *9*(1), 65–83.
- Sroufe, L. A., Egeland, B., Carlson, E. A., & Collins, W. A. (2005). *The development of the person: The Minnesota Study of Risk and Adaptation from Birth to Adulthood*. New York: Guilford.
- Taylor, C. A., Guterman, N. B., Lee, S. J., & Rathouz, P. J. (2009). Intimate partner violence, maternal stress, nativity, and risk for maternal maltreatment of young children. *American Journal of Public Health*, *99*(1), 175–183.
- Turney, K. (2012). Prevalence and correlates of stability and change in maternal depression: Evidence from the Fragile Families And Child Wellbeing Study. *PloS One*, *7*(9), e45709.
- Wang, P. S., Patrick, A., Avorn, J., Azocar, F., Ludman, E., McCulloch, J., et al. (2006). The costs and benefits of enhanced depression care to employers. *Archives of General Psychiatry*, *63*(12), 1345–1353.
- Whitaker, R. C., Orzol, S. M., & Kahn, R. S. (2006). Maternal mental health, substance use, and domestic violence in the year after delivery and subsequent behavior problems in children at age 3 years. *Archives of General Psychiatry*, *63*(5), 551–560.
- Wuslin, L. R. (2004). Is depression a major risk factor for coronary disease? A systematic review of the epidemiologic evidence. *Harvard Review of Psychiatry*, *12*, 79–93.
- U. S. Census Bureau. (2012). *Profile America facts for features*. As of July 10, 2013: http://www.census.gov/newsroom/releases/pdf/cb12ff-08_mothersday.pdf
- U.S. Department of Health and Human Services, Office of Women's Health. (2009). *Depression during and after pregnancy fact sheet*. As of February 13, 2013: <http://www.womenshealth.gov/publications/our-publications/fact-sheet/depression-pregnancy.html>
- Young, A. S., Klap, R., Sherbourne, C. D., & Wells, K. B. (2001). The quality of care for depressive and anxiety disorders in the United States. *Archives of General Psychiatry*, *58*(1), 55–61.

About the Authors

Dr. Lisa Sontag-Padilla is an Associate Behavioral/Social Scientist at RAND. Dr. Sontag-Padilla has a Ph.D. in developmental psychology from the University of Florida and completed postdoctoral training in primary care research at Cincinnati Children's Hospital Medical Center, with expertise in child development and internalizing problems. Much of her work at RAND has focused on quantitative and qualitative approaches to assessing the implementation and delivery of evidence-based interventions across multiple systems with varied processes and implementation strategies.

Dana Schultz is a Senior Policy Analyst at RAND. She has an M.P.P. from Harvard University. Her work has focused primarily on documenting implementation and assessing effectiveness of child health and well-being programs and interventions, evaluation and research design, use of quality measures, assessment of individual-level outcome changes, and rigorous approaches to program evaluation using both quantitative and qualitative methodology.

Dr. Kerry Reynolds is a Behavioral/Social Scientist at RAND. Dr. Reynolds has a Ph.D. in social/health psychology from Carnegie Mellon University, and has expertise in several areas relevant to maternal-child health, including maternal depression and parent-child decisionmaking. Her research on children and parents has spanned the developmental spectrum from infancy through emerging adulthood, and has employed both qualitative and quantitative methods.

Susan L. Lovejoy is a Health Project Associate at the RAND Corporation with experience in qualitative research and project management. Much of her work has focused on performance measurement and payment reform in various health care settings as well as program evaluation. She received her M.S. in health care policy and management from the H. John Heinz School of Public Policy and Management at Carnegie Mellon University.

Ray Firth, Policy Initiatives Director in the Office of Child Development at the University of Pittsburgh, has been fortunate to participate locally in changing Allegheny County's policy, funding, and practices for children and adults with mental illness, intellectual disabilities, and developmental challenges. Currently, he is working with local and state policymakers to utilize advances in understanding of early childhood development to implement more effective services and supports to improve the outcomes for children and families.

© Copyright 2013 RAND Corporation

www.rand.org



The RAND Corporation is a nonprofit institution that helps improve policy and decisionmaking through research and analysis. RAND focuses on the issues that matter most, such as health, education, national security, international affairs, law and business, the environment, and more. As a nonpartisan organization, RAND operates independent of political and commercial pressures. We serve the public interest by helping lawmakers reach informed decisions on the nation's pressing challenges. RAND's publications do not necessarily reflect the opinions of its research clients and sponsors. **RAND**® is a registered trademark.