



EUROPE

CHILDREN AND FAMILIES
EDUCATION AND THE ARTS
ENERGY AND ENVIRONMENT
HEALTH AND HEALTH CARE
INFRASTRUCTURE AND
TRANSPORTATION
INTERNATIONAL AFFAIRS
LAW AND BUSINESS
NATIONAL SECURITY
POPULATION AND AGING
PUBLIC SAFETY
SCIENCE AND TECHNOLOGY
TERRORISM AND
HOMELAND SECURITY

The RAND Corporation is a nonprofit institution that helps improve policy and decisionmaking through research and analysis.

This electronic document was made available from www.rand.org as a public service of the RAND Corporation.

Skip all front matter: [Jump to Page 1](#) ▼

Support RAND

[Browse Reports & Bookstore](#)

[Make a charitable contribution](#)

For More Information

Visit RAND at www.rand.org

Explore [RAND Europe](#)

View [document details](#)

Limited Electronic Distribution Rights

This document and trademark(s) contained herein are protected by law as indicated in a notice appearing later in this work. This electronic representation of RAND intellectual property is provided for non-commercial use only. Unauthorized posting of RAND electronic documents to a non-RAND Web site is prohibited. RAND electronic documents are protected under copyright law. Permission is required from RAND to reproduce, or reuse in another form, any of our research documents for commercial use. For information on reprint and linking permissions, please see [RAND Permissions](#).

This product is part of the RAND Corporation technical report series. Reports may include research findings on a specific topic that is limited in scope; present discussions of the methodology employed in research; provide literature reviews, survey instruments, modeling exercises, guidelines for practitioners and research professionals, and supporting documentation; or deliver preliminary findings. All RAND reports undergo rigorous peer review to ensure that they meet high standards for research quality and objectivity.

TECHNICAL REPORT

Postmenopausal osteoporosis management

A review of the evidence to
inform the development of
quality indicators

Annalijn Conklin, Ohid Yaqub, Claire Celia,
Ellen Nolte

Prepared for Amgen

The research described in this document was prepared for Amgen.

RAND Europe is an independent, not-for-profit research organisation whose mission is to improve policy and decision making for the public good. RAND's publications do not necessarily reflect the opinions of its research clients and sponsors.

RAND® is a registered trademark.

© Copyright 2012 RAND Corporation

Permission is given to duplicate this document for personal use only, as long as it is unaltered and complete. Copies may not be duplicated for commercial purposes. Unauthorized posting of RAND documents to a non-RAND website is prohibited. RAND documents are protected under copyright law. For information on reprint and linking permissions, please visit the RAND permissions page (<http://www.rand.org/publications/permissions.html>).

Published 2012 by the RAND Corporation
1776 Main Street, P.O. Box 2138, Santa Monica, CA 90407-2138
1200 South Hayes Street, Arlington, VA 22202-5050
4570 Fifth Avenue, Suite 600, Pittsburgh, PA 15213-2665
Westbrook Centre, Milton Road, Cambridge CB4 1YG, United Kingdom
RAND URL: <http://www.rand.org>
RAND Europe URL: <http://www.rand.org/randeurope>
To order RAND documents or to obtain additional information, contact
Distribution Services: Telephone: (310) 451-7002;
Fax: (310) 451-6915; Email: order@rand.org

Summary

Osteoporosis is the most common clinical disorder of bone metabolism. It is characterised by low bone mass and microarchitectural deterioration of bone tissue and consequent increase in bone fragility and susceptibility to fracture. Osteoporotic fractures are a major cause of morbidity; clinical complications include disability and chronic pain. It is estimated that in developed countries around 50 percent of women aged 50 and older will sustain an osteoporotic fracture during their lifetime.

Whilst there is accumulating evidence on approaches to the management of osteoporosis, the overall evidence as to the most effective interventions for the prevention and treatment of osteoporosis and associated fractures remains mixed. There is a need to draw together the available evidence to ensure high quality services are provided to those at risk of developing the condition and associated fractures.

This report aims to inform the development of quality indicators for postmenopausal osteoporosis management through (a) assessing the evidence for screening and diagnosis of osteoporosis and related risk factors, and for prevention and treatment of osteoporosis and osteoporosis-related fractures; (b) describing current practice for managing postmenopausal osteoporosis in Europe; and (c) highlighting existing gaps in the evidence base and management practices in Europe.

We have undertaken two separate sets of analyses. These involved, first, a comprehensive review of reviews as identified from the peer-reviewed and grey literature regarding the screening and diagnosis of osteoporosis and related risk factors and the prevention and treatment of osteoporosis and osteoporosis-related fractures. Secondly, we carried out case study reviews of current practices for managing postmenopausal osteoporosis in England, France, Germany and Spain, with a particular focus on the quality of care provided to those with osteoporosis and associated fractures.

There is good evidence on the effects of selected treatments on clinical outcomes of postmenopausal osteoporosis and associated fractures

Our review of reviews identified a well developed evidence base on the effects of selected treatments on clinical outcomes of postmenopausal osteoporosis and associated fractures, and on the usefulness of selected simple risk factor assessment tools to identify postmenopausal women who would benefit from further diagnostic assessment, such as dual-energy x-ray absorptiometry (DXA) measurement. We acknowledge the limitations of a review of reviews, most importantly the failure to consider more recent original studies that have not yet been included in reviews but which may be relevant and important, as for example emerging evidence examining a possible association between bisphosphonate use

and atypical fractures. Beyond this limitation, it is fair to conclude that considerable uncertainties remain in the evidence base: the optimal use (frequency, quantity, duration) of pharmacological interventions for preventive purposes; the combinations of pharmacological and/or non-pharmacological interventions that may prevent any particular type of fracture; identifying specific populations who might benefit from a given intervention (including populations who have hitherto not been studied); and the effectiveness of population-based screening. In spite of these uncertainties, the available evidence does provide some basis to inform quality improvement in clinical practice, underpinning the development of clinical guidelines in many settings in Europe.

The evidence on current approaches and practices to managing postmenopausal osteoporosis in England, France, Germany and Spain is complex

All four countries under review have introduced national-level guidelines for the management of osteoporosis. The extent to which these are implemented and/or adhered to varies, determined, in great part, by whether diagnosis and/or treatment is being reimbursed under the statutory system as well as awareness of the guidance among professionals concerned with the management of the condition.

In all countries reviewed here, there is evidence of under-diagnosis and of under-treatment, although this varies among countries, with Germany at one end of the spectrum with relatively low treatment rates (between one-fifth and one-quarter of potentially eligible women receiving treatment), and Spain at the other. Available literature points to a number of challenges faced by practitioners to implement guidance. These include lack of awareness of and knowledge about the condition, and of understanding of reimbursement mechanisms; uncertainty about responsibilities for management among providers; and restricted access to diagnostic equipment.

Improved information on reimbursement modalities and clarification of responsibilities for the management of osteoporosis and associated fractures and communication between sectors are likely to go some way to enable a more systematic approach to addressing the related societal burden in European populations.

There is considerable need for the better understanding of current approaches and practices to managing postmenopausal osteoporosis

Observations made in this study had to draw, to considerable extent, on a rather patchy evidence base, often relying on studies of small samples and/or single providers and with little systematic data collection. Furthermore, evidence that is available frequently relates to data collected in the early 2000s, so findings reported here have to be interpreted with caution.

We have identified a particular need for the establishment of routine monitoring systems to enable better understanding of contemporary patterns and trends and identify care gaps in the management of postmenopausal osteoporosis and associated fractures. Such analyses are crucial to inform targeted strategies and policies to effectively address the burden of osteoporosis and associated fractures, which is sizable and set to increase across Europe.

The systematic use of quality indicators can provide a means to enable tracking care quality. We set out considerations as a starting point for the further development of quality measures for postmenopausal osteoporosis in Europe. Such development might be able to draw, to considerable extent, on experiences in the United States, where a small set of

indicators related to the testing and the management of osteoporosis in women are already being used routinely to monitor the quality of care provided to older patients.