

# WORKING P A P E R

---

## The Prevalence and Overlap of Interstitial Cystitis/Bladder Pain Syndrome and Chronic Prostatitis/Chronic Pelvic Pain Syndrome in Men: Results of the RAND Interstitial Cystitis Epidemiology (RICE) Male Study

### Supplementary Table

ANNE M. SUSKIND, SANDRA H. BERRY,  
BRETT EWING, MARC N. ELLIOTT,  
MARIKA SUTTORP, J. QUENTIN CLEMENS

WR-959-NIDDK

September 2012

This product is part of the RAND Health working paper series. RAND working papers are intended to share researchers' latest findings and to solicit informal peer review. They have been approved for circulation by RAND Health but have not been formally edited or peer reviewed. Unless otherwise indicated, working papers can be quoted and cited without permission of the author, provided the source is clearly referred to as a working paper. RAND's publications do not necessarily reflect the opinions of its research clients and sponsors. RAND® is a registered trademark.

RICE high sensitivity and high specificity case definitions for IC/BPS.

	Female	Male
Exclusion Criteria*	Bladder cancer, urethral diverticulum, spinal cord injury, stroke, Parkinson's disease, multiple sclerosis, spina bifida, cyclophosphamide treatment, radiation treatment to pelvic area, tuberculosis affecting the bladder, genital herpes	
	Uterine cancer, ovarian cancer, vaginal cancer, pregnancy	Prostate cancer
High Sensitivity Definition	Pain, pressure, or discomfort in the pelvic area And Daytime urinary frequency 10+ or urgency due to pain, pressure, or discomfort, not fear of wetting.	
	(81% sensitivity, 54% specificity)	
High Specificity Definition	High Sensitivity Definition plus pain worsens as bladder fills And Symptoms did not resolve after treatment with antibiotics	
	And No treatment with hormone injection therapy for endometriosis  (48% sensitivity, 83% specificity)	

\*Patients were asked about exclusion criteria prior to being asked about inclusion criteria.